Author Index

Abraham, M. H. and Whiting, G. S.
Hydrogen bonding. XXI. Solvation parameters for alkylaromatic hydrocarbons from gas–liquid chromatographic data 594(1992)229
André, G., see Chauret, N. 594(1992)179
Anigbogu, V. C., Muñoz de la Peña, A., Ndou, T. T. and Warner, I. M.
Retention behavior of β-cyclodextrin complexes of anthracene and pyrene using reversed-phase liquid chromatography. The effects of *tert.*-butanol and cyclopentanol 594(1992)37
Annette, D., see Graham, W. D. 594(1992)187
Araki, T., see Kimata, K. 594(1992)87
Archambault, J., see Chauret, N. 594(1992)179
Aronson, A. L., see Tyczkowska, K. L. 594(1992)195
Auriola, S. O. K., Lepistö, A.-M., Naaranlahti, T. and Lapinjoki, S. P.
Determination of taxol by high-performance liquid chromatography–thermospray mass spectrometry 594(1992)153
Baarslag, P. C., see De Bokx, P. K. 594(1992)9
Barrett, D., see Buck, C. F. 594(1992)225
Berger, R., see Tröltzsch, C. 594(1992)111
Berger, T. A. and Deye, J. F.
Correlation between column surface area and retention of polar solutes in packed-column supercritical fluid chromatography 594(1992)291
Betto, P., Gabriele, R. and Galeffi, C.
Determination of the norlignan glucosides of Hypoxidaceae by high-performance liquid chromatography 594(1992)131
Biggs, W. R., see Jinno, K. 594(1992)105
Bogoni, P., see Zonta, F. 594(1992)137
Bohm, B. A., see Koupai-Abyazani, M. R. 594(1992)117
Bradshaw, J. S., see Johnson, D. F. 594(1992)283
Buck, C. F., Mayewski, P. A., Spencer, M. J., Whitlow, S., Twickler, M. S. and Barrett, D.
Determination of major ions in snow and ice cores by ion chromatography 594(1992)225
Burford, M. D., see Langenfeld, J. J. 594(1992)297
Caruel, H., Phemius, P., Rigal, L. and Gaset, A.
Chromatographic separation of polyols by ligand exchange. Effects of the ion-exchange resin cross-linking and size 594(1992)125
Caslavska, J. and Thormann, W.
Continuous separation and purification of proteins by recycling isotachophoresis 594(1992)361
Chalom, J., see Tod, M. 594(1992)386
Chauret, N., Côté, J., Archambault, J. and André, G.
High-performance gel permeation chromatographic analysis of immunoglobulin M produced by hybridoma cell culture 594(1992)179
Cheng, J., see Wu, C. 594(1992)243
Chloupek, R. C., Hancock, W. S. and Snyder, L. R.
Computer simulation as a tool for the rapid optimization of the high-performance liquid chromatographic separation of a tryptic digest of human growth hormone 594(1992)65
Cisowski, W., see Makuch, B. 594(1992)145
Coenegracht, P. M. J., see Wieling, J. 594(1992)45
Côté, J., see Chauret, N. 594(1992)179
De Bokx, P. K., Baarslag, P. C. and Urbach, H. P.
Modelling of displacement chromatography using non-ideal isotherms 594(1992)9
De Castro, A., see Montaño, A. 594(1992)259
Delgado, C., Sancho, P., Mendieta, J. and Luque, J.
Ligand–receptor interactions in affinity cell partitioning. Studies with transferrin covalently linked to monomethoxypoly(ethylene glycol) and rat reticulocytes 594(1992)97
De Ligny, C. L., see Gelsema, W. J. 594(1992)217
Demina, N., see Polanuer, B. 594(1992)173
Deye, J. F., see Berger, T. A. 594(1992)291
DiNunzio, J. E. and Gadde, R. R.
Development of stability indicating assay methods for the determination of hydroxamic acids in topical formulations 594(1992)209
Donnelly, J. R. and Sovocool, G. W.
Gas chromatographic elution patterns of chlorinated dioxins *versus* column polarity 594(1992)269
Doornbos, D. A., see Wieling, J. 594(1992)45
Ebermann, R., see Pichorner, H. 594(1992)400
Eguchi, M., see Johnson, D. F. 594(1992)283
Farinotti, R., see Tod, M. 594(1992)386
Feldman, B. J., Mogadeddi, H. and Osterloh, J. D.
Determination of Pb^{2+} in water by isotope dilution gas chromatography–mass spectrometry of tetraethyllead formed by reaction with sodium tetraethylborate 594(1992)275
Fetzer, J. C., see Jinno, K. 594(1992)105
Foley, J. P., see Jeansonne, M. S. 594(1992)1
Gabriele, R., see Betto, P. 594(1992)131
Gadde, R. R., see DiNunzio, J. E. 594(1992)209
Galeffi, C., see Betto, P. 594(1992)131
Gao, W., see Wu, C. 594(1992)243
Gaset, A., see Caruel, H. 594(1992)125
Gazda, K., see Makuch, B. 594(1992)145
Gelsema, W. J., De Ligny, C. L., Huigen, Y. M. and Schuring, J.
Determination of the mean ionic charge of the components of three ^{99m}Tc bone scanning agents by gel chromatography with two eluents of different electrolyte composition 594(1992)217
Gong, S., see Wu, C. 594(1992)243
Graham, W. D. and Annette, D.
Determination of ascorbic and dehydroascorbic acid in potatoes (*Solanum tuberosum*) and strawberries using ion-exclusion chromatography 594(1992)187

Author Index

Abraham, M. H. and Whiting, G. S.
Hydrogen bonding. XXI. Solvation parameters for alkylaromatic hydrocarbons from gas–liquid chromatographic data 594(1992)229
André, G., see Chauret, N. 594(1992)179
Anigbogu, V. C., Muñoz de la Peña, A., Ndou, T. T. and Warner, I. M.
Retention behavior of β-cyclodextrin complexes of anthracene and pyrene using reversed-phase liquid chromatography. The effects of *tert.*-butanol and cyclopentanol 594(1992)37
Annette, D., see Graham, W. D. 594(1992)187
Araki, T., see Kimata, K. 594(1992)87
Archambault, J., see Chauret, N. 594(1992)179
Aronson, A. L., see Tyczkowska, K. L. 594(1992)195
Auriola, S. O. K., Lepistö, A.-M., Naaranlahti, T. and Lapinjoki, S. P.
Determination of taxol by high-performance liquid chromatography–thermospray mass spectrometry 594(1992)153
Baarslag, P. C., see De Bokx, P. K. 594(1992)9
Barrett, D., see Buck, C. F. 594(1992)225
Berger, R., see Tröltzsch, C. 594(1992)111
Berger, T. A. and Deye, J. F.
Correlation between column surface area and retention of polar solutes in packed-column supercritical fluid chromatography 594(1992)291
Betto, P., Gabriele, R. and Galeffi, C.
Determination of the norlignan glucosides of Hypoxidaceae by high-performance liquid chromatography 594(1992)131
Biggs, W. R., see Jinno, K. 594(1992)105
Bogoni, P., see Zonta, F. 594(1992)137
Bohm, B. A., see Koupai-Abyazani, M. R. 594(1992)117
Bradshaw, J. S., see Johnson, D. F. 594(1992)283
Buck, C. F., Mayewski, P. A., Spencer, M. J., Whitlow, S., Twickler, M. S. and Barrett, D.
Determination of major ions in snow and ice cores by ion chromatography 594(1992)225
Burford, M. D., see Langenfeld, J. J. 594(1992)297
Caruel, H., Phemius, P., Rigal, L. and Gaset, A.
Chromatographic separation of polyols by ligand exchange. Effects of the ion-exchange resin cross-linking and size 594(1992)125
Caslavska, J. and Thormann, W.
Continuous separation and purification of proteins by recycling isotachophoresis 594(1992)361
Chalom, J., see Tod, M. 594(1992)386
Chauret, N., Côté, J., Archambault, J. and André, G.
High-performance gel permeation chromatographic analysis of immunoglobulin M produced by hybridoma cell culture 594(1992)179
Cheng, J., see Wu, C. 594(1992)243
Chloupek, R. C., Hancock, W. S. and Snyder, L. R.
Computer simulation as a tool for the rapid optimization of the high-performance liquid chromatographic separation of a tryptic digest of human growth hormone 594(1992)65
Cisowski, W., see Makuch, B. 594(1992)145
Coenegracht, P. M. J., see Wieling, J. 594(1992)45
Côté, J., see Chauret, N. 594(1992)179
De Bokx, P. K., Baarslag, P. C. and Urbach, H. P.
Modelling of displacement chromatography using non-ideal isotherms 594(1992)9
De Castro, A., see Montaño, A. 594(1992)259
Delgado, C., Sancho, P., Mendieta, J. and Luque, J.
Ligand–receptor interactions in affinity cell partitioning. Studies with transferrin covalently linked to monomethoxypoly(ethylene glycol) and rat reticulocytes 594(1992)97
De Ligny, C. L., see Gelsema, W. J. 594(1992)217
Demina, N., see Polanuer, B. 594(1992)173
Deye, J. F., see Berger, T. A. 594(1992)291
DiNunzio, J. E. and Gadde, R. R.
Development of stability indicating assay methods for the determination of hydroxamic acids in topical formulations 594(1992)209
Donnelly, J. R. and Sovocool, G. W.
Gas chromatographic elution patterns of chlorinated dioxins *versus* column polarity 594(1992)269
Doornbos, D. A., see Wieling, J. 594(1992)45
Ebermann, R., see Pichorner, H. 594(1992)400
Eguchi, M., see Johnson, D. F. 594(1992)283
Farinotti, R., see Tod, M. 594(1992)386
Feldman, B. J., Mogadeddi, H. and Osterloh, J. D.
Determination of Pb^{2+} in water by isotope dilution gas chromatography–mass spectrometry of tetraethyllead formed by reaction with sodium tetraethylborate 594(1992)275
Fetzer, J. C., see Jinno, K. 594(1992)105
Foley, J. P., see Jeansonne, M. S. 594(1992)1
Gabriele, R., see Betto, P. 594(1992)131
Gadde, R. R., see DiNunzio, J. E. 594(1992)209
Galeffi, C., see Betto, P. 594(1992)131
Gao, W., see Wu, C. 594(1992)243
Gaset, A., see Caruel, H. 594(1992)125
Gazda, K., see Makuch, B. 594(1992)145
Gelsema, W. J., De Ligny, C. L., Huigen, Y. M. and Schuring, J.
Determination of the mean ionic charge of the components of three ^{99m}Tc bone scanning agents by gel chromatography with two eluents of different electrolyte composition 594(1992)217
Gong, S., see Wu, C. 594(1992)243
Graham, W. D. and Annette, D.
Determination of ascorbic and dehydroascorbic acid in potatoes (*Solanum tuberosum*) and strawberries using ion-exclusion chromatography 594(1992)187

Grateful, T. M. and Lightfoot, Jr., E. N.
Finite difference modelling of continuous-flow electrophoresis 594(1992)341
Grunau, J. A. and Swiader, J. M.
Chromatography of 99 amino acids and other ninhydrin-reactive compounds in the Pickering lithium gradient system 594(1992)165
Hancock, W. S., see Chloupek, R. C. 594(1992)65
Hartwick, R. A., see Wang, T. 594(1992)325
Hawthorne, S. B., see Langenfeld, J. J. 594(1992)297
Hirokawa, T., see Hu, J. 594(1992)371
Hodges, R. S., see Zhu, B.-Y. 594(1992)75
Hu, J., Hirokawa, T., Nishiyama, F., Kiso, Y., Ito, K. and Shoto, E.
Study of isotachophoretic separation behaviour of metal cations by means of particle-induced X-ray emission. III. Analysis of a crude rare earth chloride from monazite 594(1992)371
Huigen, Y. M., see Gelsema, W. J. 594(1992)217
Ito, K., see Hu, J. 594(1992)371
Itoh, K., see Jinno, K. 594(1992)105
Jeansonne, M. S. and Foley, J. P.
Improved equations for the calculation of chromatographic figures of merit for ideal and skewed chromatographic peaks 594(1992)1
Jenke, D. R.
Determination of solute–polymer interaction constants via chromatography with polymer capillaries 594(1992)29
Jessner, G., see Pichorner, H. 594(1992)400
Jinno, K., Yamamoto, K., Ueda, T., Nagashima, H., Itoh, K., Fetzer, J. C. and Biggs, W. R.
Liquid chromatographic separation of all-carbon molecules C_{60} and C_{70} with multi-legged phenyl group bonded silica phases 594(1992)105
Johnson, D. F., Bradshaw, J. S., Eguchi, M., Rossiter, B. E., Lee, M. L., Petersson, P. and Markides, K. E.
Synthesis of (1*R-trans*)-N,N′-1,2-cyclohexylenebisbenzamideoligodimethylsiloxane copolymers for use as chiral stationary phases for capillary supercritical fluid chromatography 594(1992)283
Jonkman, J. H. G., see Wieling, J. 594(1992)45
Kakemoto, M.
Simultaneous determination of sorbic acid, dehydroacetic acid and benzoic acid by gas chromatography–mass spectrometry 594(1992)253
Khuhawar, M. Y., see Sarafraz-Yazdi, A. 594(1992)395
Kimata, K., Tanaka, N. and Araki, T.
Suppression of the effect of metal impurities in alkylsilylated silica packing materials 594(1992)87
Kiso, Y., see Hu, J. 594(1992)371
Korori, S. A. A., see Pichorner, H. 594(1992)400
Kostenko, V. G., see Nemirovskii, V. D. 594(1992)23
Koupai-Abyazani, M. R., McCallum, J. and Bohm, B. A.
Identification of the constituent flavanoid units in sainfoin proanthocyanidins by reversed-phase high-performance liquid chromatography 594(1992)117
Kouwatli, H., see Tod, M. 594(1992)386
Langenfeld, J. J., Burford, M. D., Hawthorne, S. B. and Miller, D. J.
Effects of collection solvent parameters and extraction cell geometry on supercritical fluid extraction efficiencies 594(1992)297
Lapinjoki, S. P., see Auriola, S. O. K. 594(1992)153
Lee, H.-B. and Peart, T. E.
Supercritical carbon dioxide extraction of resin and fatty acids from sediments at pulp mill sites 594(1992)309
Lee, H. K., see Yeo, S. K. 594(1992)335
Lee, M. L., see Johnson, D. F. 594(1992)283
Legendre, J.-Y., see Tod, M. 594(1992)386
Lepistö, A.-M., see Auriola, S. O. K. 594(1992)153
Li, S. F. Y., see Yeo, S. K. 594(1992)335
Lightfoot, Jr., E. N., see Grateful, T. M. 594(1992)341
Little, R. J., see Tsuji, K. 594(1992)317
Loew, S., see Pichorner, H. 594(1992)400
Lu, X., see Wu, C. 594(1992)243
Luque, J., see Delgado, C. 594(1992)97
Mahuzier, G., see Tod, M. 594(1992)386
Makuch, B., Gazda, K. and Cisowski, W.
Isolation of aristolochic acids HA_3I and HA_3II by preparative liquid chromatography 594(1992)145
Mant, C. T., see Zhu, B.-Y. 594(1992)75
Markides, K. E., see Johnson, D. F. 594(1992)283
Maruyama, M. and Nagayoshi, T.
Reversed-phase liquid chromatographic behaviour of alkylamines with amperometric detection and its application to trace analysis 594(1992)159
Masotti, P., see Zonta, F. 594(1992)137
Mayewski, P. A., see Buck, C. F. 594(1992)225
McCallum, J., see Koupai-Abyazani, M. R. 594(1992)117
Mendieta, J., see Delgado, C. 594(1992)97
Mensink, C. K., see Wieling, J. 594(1992)45
Michalik, M., see Tröltzsch, C. 594(1992)111
Miller, D. J., see Langenfeld, J. J. 594(1992)297
Misra, S., see Sharma, S. D. 594(1992)379
Mogadeddi, H., see Feldman, B. J. 594(1992)275
Monakhova, N. I., see Nemirovskii, V. D. 594(1992)23
Montaño, A., De Castro, A., Rejano, L. and Sánchez, A.-H.
Analysis of zapatera olives by gas and high-performance liquid chromatography 594(1992)259
Muñoz de la Peña, A., see Anigbogu, V. C. 594(1992)37
Naaranlahti, T., see Auriola, S. O. K. 594(1992)153
Nagashima, H., see Jinno, K. 594(1992)105
Nagayoshi, T., see Maruyama, M. 594(1992)159
Nambara, T.
Advances in steroid analysis (edited by S. Görög) (Book Review) 594(1992)403
Ndou, T.T., see Anigbogu, V.C. 594(1992)37
Nemirovskii, V. D., Monakhova, N. I. and Kostenko, V. G.
High-performance liquid chromatographic study of the correlation between the physico-chemical parameters of furan and benzene aldehydes and the rate of their metabolism by yeast 594(1992)23
Nishiyama, F., see Hu, J. 594(1992)371
Olsen, B. A., Stafford, J. D. and Reed, D. E.
Determination of dirithromycin purity and related substances by high-performance liquid chromatography 594(1992)203

Osterloh, J. D., see Feldman, B. J. 594(1992)275
Peart, T. E., see Lee, H.-B. 594(1992)309
Petersson, P., see Johnson, D. F. 594(1992)283
Phemius, P., see Caruel, H. 594(1992)125
Pichorner, H., Loew, S., Korori, S. A. A., Jessner, G. and Ebermann, R.
Fast and sensitive staining technique for catalase in polyacrylamide gel 594(1992)400
Polanuer, B., Sholin, A., Demina, N. and Rumiantseva, N.
Determination of glutamine, glutamic acid and pyroglutamic acids using high-performance liquid chromatography on dynamically modified silica 594(1992)173
Poulou, M., see Tod, M. 594(1992)386
Reed, D. E., see Olsen, B. A. 594(1992)203
Rejano, L., see Montaño, A. 594(1992)259
Rigal, L., see Caruel, H. 594(1992)125
Rossiter, B. E., see Johnson, D. F. 594(1992)283
Roux-de Balmann, H. and Sanchez, V.
Continuous-flow electrophoresis: a separation criterion applied to the separation of model proteins 594(1992)351
Rumiantseva, N., see Polanuer, B. 594(1992)173
Sánchez, A.-H., see Montaño, A. 594(1992)259
Sanchez, V., see Roux-de Balmann, H. 594(1992)351
Sancho, P., see Delgado, C. 594(1992)97
Sarafraz-Yazdi, A., Khuhawar, M. Y. and Uden, P. C.
Gas chromatography of chloride and bromide as phenylboronic acid/mercuric nitrate derivatives with microwave induced plasma atomic emission detection 594(1992)395
Schuring, J., see Gelsema, W. J. 594(1992)217
Sharma, S. D. and Misra, S.
Quantitative separation of Cr^{3+} from Mo^{6+}, W^{6+}, Hg_2^{2+}, Cu^{2+} and Pb^{2+}. Chromatographic behaviour of 51 cations on papers impregnated with Sn(IV)-based inorganic ion exchangers in complex-forming acid systems 594(1992)379
Sholin, A., see Polanuer, B. 594(1992)173
Shoto, E., see Hu, J. 594(1992)371
Snyder, L. R., see Chloupek, R. C. 594(1992)65
Sovocool, G. W., see Donnelly, J. R. 594(1992)269
Spencer, M. J., see Buck, C. F. 594(1992)225
Stafford, J. D., see Olsen, B. A. 594(1992)203
Stancher, B., see Zonta, F. 594(1992)137
Swiader, J. M., see Grunau, J. A. 594(1992)165
Tanaka, N., see Kimata, K. 594(1992)87
Thormann, W., see Caslavska, J. 594(1992)361
Tod, M., Legendre, J.-Y., Chalom, J., Kouwatli, H., Poulou, M., Farinotti, R. and Mahuzier, G.
Primary and secondary amine derivatization with luminarins 1 and 2: separation by liquid chromatography with peroxyoxalate chemiluminescence detection 594(1992)386
Tröltzsch, C., Berger, R. and Michalik, M.
Liquid chromatographic isolation and structural elucidation of methoxymethylene dimethyl dodecatrienes catalytically synthesized from methanol and three moles of isoprene 594(1992)111
Tsuji, K. and Little, R. J.
Charge-reversed, polymer-coated capillary column for the analysis of a recombinant chimeric glycoprotein 594(1992)317
Twickler, M. S., see Buck, C. F. 594(1992)225
Tyczkowska, K. L., Voyksner, R. D. and Aronson, A. L.
Solvent degradation of cloxacillin *in vitro*. Tentative identification of degradation products using thermospray liquid chromatography–mass spectrometry 594(1992)195
Uden, P. C., see Sarafraz-Yazdi, A. 594(1992)395
Ueda, T., see Jinno, K. 594(1992)105
Urbach, H. P., see De Bokx, P. K. 594(1992)9
Voyksner, R. D., see Tyczkowska, K. L. 594(1992)195
Wang, T. and Hartwick, R. A.
Capillary modification and evaluation using streaming potential and frontal chromatography for protein analysis in capillary electrophoresis 594(1992)325
Warner, I. M., see Anigbogu, V. C. 594(1992)37
Watson, E. and Yao, F.
Fast protein monitoring in fermentation broth using non-porous micropellicular reversed-phase columns 594(1992)392
Whiting, G. S., see Abraham, M. H. 594(1992)229
Whitlow, S., see Buck, C. F. 594(1992)225
Wieling, J., Coenegracht, P. M. J., Mensink, C. K., Jonkman, J. H. G. and Doornbos, D. A.
Selection of robust combinations of extraction liquid composition and internal standard. Monte Carlo simulation of improvement of assay methods with liquid–liquid extraction prior to high-performance liquid chromatography 594(1992)45
Wu, C., Cheng, J., Gao, W., Zeng, Z., Lu, X. and Gong, S.
Preparation and capillary column gas chromatographic characterization of aza-crown ether polysiloxane stationary phases 594(1992)243
Yamamoto, K., see Jinno, K. 594(1992)105
Yao, F., see Watson, E. 594(1992)392
Yeo, S. K., Lee, H. K. and Li, S. F. Y.
Separation of plant growth regulators by capillary electrophoresis 594(1992)335
Zeng, Z., see Wu, C. 594(1992)243
Zhu, B.-Y., Mant, C. T. and Hodges, R. S.
Mixed-mode hydrophilic and ionic interaction chromatography rivals reversed-phase liquid chromatography for the separation of peptides 594(1992)75
Zonta, F., Stancher, B., Bogoni, P. and Masotti, P.
High-performance liquid chromatography of okadaic acid and free fatty acids in mussels 594(1992)137